

# Multilinear Hirota operators and integrability

Il'in I.A.<sup>a,b</sup>, Noshchenko D.S.<sup>a,b</sup>, Perezhogin A.S.<sup>a,b</sup>

<sup>a</sup>*Institute of Cosmophysical Research and Radio Wave Propagation of the Far Eastern Branch of Russian Academy of Science, 684034 Kamchatka region, Elizovskiy district, Paratunka, Mirnaya str., 7.*

<sup>b</sup>*Vitus Bering Kamchatka State University, 683032, Kamchatka region, Elizovskiy district, Petropavlovsk-Kamchatsky, Pogranichnaya str., 4.*

## 1. Introduction

Hirota's bilinear derivative [1, 4, 5, 6] is a powerful tool for construction of integrable nonlinear equations. For instance, the KdV equation

$$u_t + 6uu_x + u_{xxx} = 0, u(x, t) = \frac{\partial^2}{\partial x^2} \log f(x, t) \quad (1)$$

transforms to bilinear form

$$D_x(D_x^3 + D_t) f \cdot f = 0, \quad (2)$$

where

$$D_x^m D_t^n = (\partial_x - \partial_{x'})^m (\partial_t - \partial_{t'})^n f(x, t) \cdot f(x', t')|_{x'=x, t'=t} \quad (3)$$

or, equivalently,

$$D_x^m f \cdot f = \frac{d^m}{dy^m} f(x-y) f(x+y) \Big|_{y=0} \quad (4)$$

Any bilinear equation  $G(D_x, D_t) f \cdot f = 0$ , where  $G$  is a polynomial, has at least 2SS. But 3SS condition is much more stronger. For instance, among the equations

$$(D_x D_t + D_x^n) f \cdot f = 0 \quad (5)$$

only  $n = 4, n = 6$  has 3SS (KdV and SK equations). We name  $n = 7$  as “new” and  $n = 9$  as “tail” equations.

Here are some integrable bilinear equations, listed in [4]:

$$(D_x^4 - 4D_x D_t + 3D_y^2) f \cdot f = 0, \quad (6)$$

$$(D_x^3 D_t + aD_x^2 + D_t D_y) f \cdot f = 0, \quad (7)$$

$$(D_x^4 - D_x D_t^3 + aD_x^2 + bD_x D_t + cD_t^2) f \cdot f = 0, \quad (8)$$

$$(D_x^6 + 5D_x^3 D_t - 5D_t^2 + D_x D_y) f \cdot f = 0. \quad (9)$$

## 2. Multilinear Hirota operators

Consider k-linear gauge invariant Hirota operator, introduced in [4], [1]

$$(H_{k,l}^m)_x = (\partial_{x_1} + \gamma_1 \partial_{x_2} + \dots + \gamma_{k-1} \partial_{x_k})^m f(x_1) \cdot \dots \cdot f(x_k)|_{x_i=x, i=1\dots k} \quad (10)$$

$\gamma$  - primitive kth root of unity. For  $k = 3$  we have

$$T^3 = H_{3,1} = \partial_1 + e^{\frac{2}{3}i\pi} \partial_2 + e^{\frac{4}{3}i\pi} \partial_3 \quad (11)$$

$$T^{*3} = H_{3,2} = \partial_1 + e^{\frac{4}{3}i\pi} \partial_2 + e^{\frac{2}{3}i\pi} \partial_3 \quad (12)$$

For example, Chazy 3rd order integrable equation can be written as a single trilinear determinant or as a pair of bilinear equations ??.

$$\begin{vmatrix} f'' & f' & f \\ f''' & f'' & f' \\ f'''' & f''' & f'' \end{vmatrix} = (T^6 - T^3 T^{*3}) f \cdot f \cdot f = \quad (13)$$

$$= f (D_x^3 (D_{x'} + D_x^3) f \cdot f), \quad (14)$$

with  $D_x (D_{x'} + D_x^3) f \cdot f = 0$ .

Denote  $Q_1^l Q_2^m Q_3^n = Q^{l,m,n}$  - quadrilinear Hirota operator. In a similar way, the 4th order determinant

$$\begin{vmatrix} f'''' & f''' & f'' & f' \\ f'''' & f'''' & f''' & f'' \\ f'''' & f'''' & f'''' & f''' \\ f'''' & f'''' & f'''' & f'''' \end{vmatrix} = \quad (15)$$

$$\frac{1}{5 \cdot 7 \cdot 2^{14}} (45 Q^{9,2,1} + 38 Q^{7,2,3} - 7 Q^{5,2,5} - 8 Q^{8,4,0} + 72 Q^{2,8,2} - 140 Q^{3,6,3}), \quad (16)$$

which is a weighted sum of the following operator determinants

$$\begin{vmatrix} Q_1^9 & 0 & Q_1^5 \\ 0 & Q_2^2 & 0 \\ Q_3^5 & 0 & Q_3 \end{vmatrix}, \begin{vmatrix} 0 & Q_1^8 & Q_1^2 \\ Q_2^8 & 0 & Q_2^4 \\ -1 & Q_3^2 & 0 \end{vmatrix}, \begin{vmatrix} Q_1^7 & Q_1^3 & Q_1^5 \\ Q_2^6 & Q_2^2 & 0 \\ Q_3^5 & 0 & Q_3^3 \end{vmatrix} \quad (17)$$

Sometimes it is helpful to use the generating series for multilinear Hirota operators:

$$H_k f \cdot \dots \cdot f = H \left[ \prod_{j=1}^k f_j \left( x + \sum_{m=1}^{k-1} y_m e^{jm \frac{2\pi i}{k}} \right) \right] \Big|_{y_1=\dots=y_{k-1}=0} \quad (18)$$

Quadrilinear operator can be derived as follows:

$$Q^{l,m,n} = \frac{\partial^{l+j+k}}{\partial y_1^l \partial y_2^m \partial y_3^n} (f(x + y_1 + y_2 + y_3) f(x + iy_1 - y_2 - iy_3) \cdot \quad (19)$$

$$\cdot f(x - y_1 + y_2 - y_3) f(x - iy_1 - y_2 + iy_3))|_{y_1=y_2=y_3=0}$$

Why roots of unity? Consider Taylor expansion of the product

$$f(x + c_1 y) f(x + c_2 y) f(x + c_3 y)|_{y_1=y_2=y_3=0} \quad (20)$$

Expanding the coefficients, we get

$$c_1 = (c_3 + c_2 + c_1) f^2 f_x, \quad (21)$$

$$c_2 = ((c_2 + c_1) c_3 + c_1 c_2) f f_x^2 + 1/2 (c_3^2 + c_2^2 + c_1^2) f^2 f_{x,x}. \quad (22)$$

The coefficients are clearly homogeneous of weight  $kp + s$  and satisfy the following property:

$$c_s(e^x \cdot f) = c_s(f) + \sum_{k=0}^{s-1} a_k c_k(f) \quad (23)$$

Equalling  $c_1$  to zero implies all other coeffs to be gauge-invariant. When both of  $c_1, c_2$  are zero, only powers by modulo 3 left in expansion, and we immediately get  $\{T^i\}$  – sequence of trilinear operators.

Identically zero condition:

$$\sum_{i=1}^{k-1} c_i \cdot \text{binomial}(k-2, i) m_i \bmod k \neq 0 \quad (24)$$

implies  $H_{n,1}^{i_1} H_{n,2}^{i_2} \dots H_{n,n-1}^{i_{n-1}} \equiv 0$  ( $c_i = 1$  if  $k$  is prime). Example:  $i - j \bmod 3 = 0, i + 2j - k \bmod 4 = 0$ .

Invariance condition:

$$(H_k(e^{x^{i-1}} f))^m = H_k^m f, 0 < i \leq m, \quad i = m \Leftrightarrow m|k$$

Now we can obtain the  $\varepsilon$ -expansion of  $k$ -linear Hitora operator, which is useful for finding solitonic or polynomial solutions. We use the differentiation rule

$$H_k(f \cdot f \cdot \dots \cdot 1) = \frac{d}{df} H_k(f \cdot f \cdot \dots \cdot f)$$

For simplicity, we consider  $k = 3$ .

$$\mathcal{T} = \sum_{i,j} T^i T^{*j}, \quad (25)$$

is any finite linear combination of  $T$ 's. Substituting  $f = \sum_i \varepsilon^i f_i$  in 25 ( $\varepsilon$  is a formal parameter), we get the following perturbation expansion:

$$\begin{aligned}
\varepsilon & L f_1 \\
\varepsilon^2 & (\mathcal{T} f_1 \cdot f_1 \cdot f_1)'_{f_1} + L f_2 \\
\varepsilon^3 & (\mathcal{T} f_1 \cdot f_1 \cdot f_1) + \sum_{i=1}^n (\mathcal{T} f_1 \cdot f_1 \cdot f_1)''_{f_1^{(0)} f_1^{(1)}} f_2^{(i)} + L f_3 \\
\varepsilon^4 & \sum_{i=1}^n (\mathcal{T} f_1 \cdot f_1 \cdot f_1)'_{f_1^{(1)}} f_2^{(i)} + \sum_{i=1}^n (\mathcal{T} f_1 \cdot f_1 \cdot f_1)''_{f_1^{(0)} f_1^{(1)}} f_3^{(i)} + L f_4 \\
\varepsilon^5 & \sum_{i=1}^n \sum_{j=1}^n (\mathcal{T} f_1 \cdot f_1 \cdot f_1)''_{f_1^{(i)} f_1^{(j)}} f_3^{(i)} f_3^{(j)} + \sum_{i=1}^n (\mathcal{T} f_1 \cdot f_1 \cdot f_1)'_{f_1^{(1)}} f_3^{(i)} + \\
& + \sum_{i=1}^n (\mathcal{T} f_1 \cdot f_1 \cdot f_1)''_{f_1^{(0)} f_1^{(1)}} f_4^{(i)} + L f_5 \\
& \vdots
\end{aligned}$$

The following table shows multilinear representation of some known equations:

| Equation | Bilinear           | Trilinear                                                                      | Quadrilinear                                                                                                                              |
|----------|--------------------|--------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| KdV 3    | $D_x(D_x^3 + D_t)$ | $2T_x^2 T_x^{*2} + 3T_x^* T_t$                                                 | $2G1_x^2 G3_x^2 + G1_x^2 G2_t^1$                                                                                                          |
| KdV 5    | -                  | $T_x^4 T_x^* + T_x^* T_t$<br>$20T_x^3 T_x^{*3} + 7T_x^6 + 27T_x^* T_t$         | $G2_x^1 G3_x^6 + 5G1_x^1 G2_x^3 G3_x^3 + 12G1_x^2 G2_t^1$                                                                                 |
| SK 5     | $D_x(D_x^5 + D_t)$ | $10T_x^3 T_x^{*3} - T_x^6 + 9T_x^* T_t$                                        |                                                                                                                                           |
| SK 7     |                    | $T_x^5 T_x^{*2} + T_x^2 T_t$<br>$14T_x^4 T_x^{*4} + 14T_x^7 T_x^* + 27T_x T_t$ | $3G3_x^8 + 7G1_x^2 G3_x^6 - 42G1_x^2 G2_x^4 G3_x^2 + 64G1_x^2 G2_t^1$<br>$7G1_x^2 G2_x^3 G3_x^4 + G1_x^1 G2_x^1 G3_x^7 + 16G1_x^2 G2_t^1$ |
| “new” 7  | $D_x(D_x^7 + D_t)$ | $35T_x^4 T_x^{*4} - 4T_x^7 T_x^* + 27T_x T_t$                                  |                                                                                                                                           |
| “tail” 9 | $D_x(D_x^9 + D_t)$ | $14T_x^5 T_x^{*5} - 5T_x^8 T_x^{*2} + 9T_x T_t$                                | $7G2_x^4 G3_x^{*4} - G1_x G2_x^2 G3_x^5 + 12G1_x^2 G2_t^1$                                                                                |

In the next section we investigate two trilinear integrable equations, considered in [4], and we show that they are representable as systems of bilinear equations. For some information on Painleve test, Fuchs indicies and other integrability features we refer reader to [3].

### 3. J8 equation

$$\left( 4T_x^7 T_x^* + 5T_x^4 T_x^{*4} + \alpha(20T_x^3 T_x^{*3} + 7T_x^6) + 9\beta T_x^2 T_x^* T_t + \frac{9\alpha\beta}{2} T_x T_t^* \right) f \cdot f \cdot f = 0 \quad (26)$$

This equation passes Painleve test and has 3SS for arbitrary  $\alpha, \beta$ . Fuchs index

$$\begin{pmatrix} -1 & 2 & 3 & 4 & 8 & 9 & 10 \\ -2 & -1 & 3 & 5 & 8 & 10 & 12 \\ -7 & -5 & -1 & 8 & 10 & 12 & 18 \end{pmatrix} \quad (27)$$

For  $f(x) = 1 + e^{p_i x - q_i t}$  dispersion law is

$$q_i = 2 \frac{p_i^5 (3\alpha + p_i^2)}{\beta (2 p_i^2 + \alpha)} \quad (28)$$

and for 2SS  $f(x) = 1 + e^{p_1 x - q_1 t} + e^{p_2 x - q_2 t} + A_{1,2} e^{p_1 x - q_1 t} e^{p_2 x - q_2 t}$  we have

$$A_{1,2} = \frac{(p_1 - p_2)^2 (2 p_1^2 - 2 p_2 p_1 + 3 \alpha + 2 p_2^2)}{(p_1 + p_2)^2 (2 p_1^2 + 2 p_2 p_1 + 3 \alpha + 2 p_2^2)} \quad (29)$$

so  $\alpha = 0$  gives SK5 solution. Next we take  $\alpha = 0, \beta = 1$  and show that (26) can be derived from SK5.

SK5 equation:

$$(u_{x,x,x,x} + 30 u_{x,x} u + 60 u_x^3)_x + u_t = 0 \quad (30)$$

also can be written in a bilinear form with  $u = \partial_{x,x} \log(f)$ :

$$D_x(D_x^5 + D_t) f \cdot f = 0 \quad (31)$$

which is equivalent to

$$(-T_x^6 + 10 T_x^3 T_x^{*3} + 9 T_x T_t^*) f \cdot f \cdot f = 0 \quad (32)$$

Now we compare two forms:

$$J8 = T_x T_x^* (4 T_x^6 + 5 T_x^3 T_x^{*3} + 9 T_x T_t^*) f \cdot f \cdot f \quad (33)$$

and bilinear form of SK5 (31). J8 evaluates to

$$J8 = (f_{t,x,x,x} + f_{x,x,x,x,x,x,x}) f^2 + (-f_{x,x,x} f_t - 5 f_{x,x,x,x}^2 + 3 f_{x,x} f_{t,x} + 4 f_{x,x,x,x,x} f_{x,x,x} + 8 f_{x,x,x,x,x,x} f_{x,x} - 8 f_{x,x,x,x,x,x,x} f_x - 3 f_{t,x,x} f_x) f - 40 f_{x,x} f_{x,x,x}^2 + 20 f_{x,x,x,x,x,x} f_x^2 + 60 f_{x,x,x,x} f_{x,x}^2 + 20 f_x f_{x,x,x,x} f_{x,x,x} - 60 f_x f_{x,x,x,x,x} f_{x,x}$$

Mult. (31) by  $f$  and diff. twice, we get

$$(SK5 \cdot f)_{x,x} = (f_{t,x,x,x} + f_{x,x,x,x,x,x,x}) f^2 + (-2 f_{x,x,x,x,x,x,x} f_x + 5 f_{x,x,x,x,x,x} f_{x,x} + 4 f_{x,x,x,x,x} f_{x,x,x} + 3 f_{t,x,x} f_x - f_{x,x,x} f_t - 5 f_{x,x,x,x}^2) f - 10 f_{x,x} f_{x,x,x}^2 + 12 f_x f_{x,x,x,x,x} f_{x,x} - 10 f_x f_{x,x,x,x} f_{x,x,x} - 3 f_x f_{x,x} f_t - 10 f_{x,x,x,x,x,x} f_x^2 + 15 f_{x,x,x,x} f_{x,x}^2$$

Direct computation shows that

$$J8 - (SK5 \cdot f)_{x,x} = 3(SK5 \cdot f_x)_x + R(x) \quad (34)$$

where

$$R(x) = -9 (f_{x,x,x,x,x,x,x} f - 5 f_{x,x,x,x,x,x} f_x + 9 f_{x,x,x,x,x} f_{x,x} - 5 f_{x,x,x,x} f_{x,x,x} + f_{t,x,x} f - f_{x,x} f_t) f_x \quad (35)$$

which is nothing but  $(SK5)_x$  mult. by  $-9f_x$ . So

$$J8 = (SK5 \cdot f)_{x,x} + 3(SK5 \cdot f_x)_x - 9f_x(SK5)_x = f(SK5)_{x,x} - 4D_x(SK5 \cdot f) \quad (36)$$

#### 4. J10 equation

$$\left(5T_x^8T_x^{*2} + 4T_x^5T_x^{*5} + \alpha(4T_x^7T_x^* + 5T_x^4T_x^{*4}) + \beta(20T_x^3T_x^{*2}T_t^* + 7T_x^5T_t) + 6\alpha\beta T_x^2T_x^*T_t^* + \frac{3\beta^2}{2}T_tT_t^*\right) f \cdot f \cdot f = 0 \quad (37)$$

Fuchs indices:

$$\begin{pmatrix} -1 & 2 & 3 & 4 & 5 & 6 & 12 & 13 \\ -2 & -1 & 2 & 3 & 5 & 9 & 12 & 16 \\ -3 & -2 & -1 & 4 & 5 & 11 & 12 & 18 \\ -13 & -11 & -7 & -1 & 12 & 16 & 18 & 30 \end{pmatrix} \quad (38)$$

Dispersion law:

$$q_i = \frac{(2\alpha + 9p_i^2 + \pm(\sqrt{4\alpha^2 + 30\alpha p_i^2 + 75p_i^4}))p_i^3}{\beta} \quad (39)$$

Again, 2SS exists for arbitrary  $\alpha, \beta$ , but its coeff.  $A_{1,2}$  is very large. 3SS also exists. In order to exclude J8 terms from (40), we assume  $\alpha = 0$ :

$$J10 = \left(5T_x^8T_x^{*2} + 4T_x^5T_x^{*5} + \beta(20T_x^3T_x^{*2}T_t^* + 7T_x^5T_t) + \frac{3\beta^2}{2}T_tT_t^*\right) f \cdot f \cdot f = 0 \quad (40)$$

2SS for J10 is given by:

$$A_{1,2} = -\frac{(p_2^2 + \sqrt{3}p_1p_2 + p_1^2)(p_2^2 - p_1p_2 + p_1^2)(-p_2 + p_1)^2}{(-p_2^2 + \sqrt{3}p_1p_2 - p_1^2)(p_2^2 + p_1p_2 + p_1^2)(p_2 + p_1)^2} \quad (41)$$

This will lead us to associated bilinear form given by:

$$\left(q - \frac{(9 - 5\sqrt{3})p^5}{\beta}\right)\left(q - \frac{(9 + 5\sqrt{3})p^5}{\beta}\right) \quad (42)$$

Corr. bilinear operator is defined uniquely:

$$B1 = \left(\frac{6}{\beta^2}D_x^{10} + \frac{18}{\beta}D_x^5D_t + D_t^2\right) f \cdot f \quad (43)$$

This eq. (43) has 2SS, but not 3SS. Its coeff.:

$$A_{1,2} = \frac{(p_2^2 - p_1p_2 + p_1^2)(2p_1^2 - 3p_1p_2 + \sqrt{3}p_1p_2 + 2p_2^2)(p_1^2 + \sqrt{3}p_1p_2 + p_2^2)(p_1 - p_2)^2}{(-p_1^2 + \sqrt{3}p_1p_2 - p_2^2)(-2p_1^2 - 3p_1p_2 + \sqrt{3}p_1p_2 - 2p_2^2)(p_2^2 + p_1p_2 + p_1^2)(p_1 + p_2)^2} \quad (44)$$

Now we are going to represent (40) as a system of bilinear equations with (43) plus some additional relations. At first we rewrite (43) in trilinear form.

$$(f \cdot D_t^2) = \frac{2}{3} T_t T_t^* \quad (45)$$

$$(f \cdot D_x^6) = \frac{2}{27} (-T_x^6 + 10 T_x^3 T_x^{*3}) \quad (46)$$

$$(f \cdot D_x^5 D_t) = \frac{2}{27} (-T_x^5 T_t + 10 T_x^3 T_x^{*2} T_t^*) \quad (47)$$

$$(f \cdot D_x^{10}) = \frac{2}{27} (-5 T_x^8 T_x^{*2} + 14 T_x^5 T_x^{*5}) \quad (48)$$

Dividing by lead coeff., we get

$$f \cdot B1 = -5 T_x^8 T_x^{*2} + 14 T_x^5 T_x^{*5} + 3\beta (-T_x^5 T_t + 10 T_x^3 T_x^{*2} T_t^*) + \frac{3\beta^2}{2} T_t T_t^* \quad (49)$$

So thier difference is symmetrical:

$$J10 - (f \cdot B1) = 10(\beta(T_x^5 T_t - T_x^3 T_x^{*2} T_t^*) + (T_x^8 T_x^{*2} - T_x^5 T_x^{*5})) \quad (50)$$

Note here coeffs summ to zero! Here both terms  $T_x^5 T_t - T_x^3 T_x^{*2} T_t^*$ ,  $T_x^8 T_x^{*2} - T_x^5 T_x^{*5}$  are zero for  $f(x, t) = 1 + e^{p_x - q_x}$  for any  $p, q$ . Subs.  $f(x, t) = 1 + e^{p_1 x - q_1 x} + e^{p_2 x - q_2 x}$  in the second term and factorizing, we get

$$-3 p_1^2 e^{p_1 x - q_1 t + p_2 x - q_2 t} p_2^2 (-p_2 + p_1)^2 (p_2^2 - p_1 p_2 + p_1^2)^2 \quad (51)$$

This is 2SS numer for the new7 eq. One can check that

$$(T_x^8 T_x^{*2} - T_x^5 T_x^{*5}) f \cdot f \cdot f \equiv c_1 D_x^3 (D_\tau + D_x^7) f \cdot f, \quad (52)$$

with

$$D_x (D_\tau + D_x^7) f \cdot f = 0 \quad (53)$$

(it is easier with  $u(x, t, \tau)$  variable)

Another term is similar to Lax5 and it splits to

$$(T_x^5 T_t - T_x^3 T_x^{*2} T_t^*) f \cdot f \cdot f \equiv c_2 D_x^3 (D_{\tau'} + D_x^2 D_t) f \cdot f, \quad (54)$$

with

$$D_x (D_{\tau'} + D_x^2 D_t) f \cdot f = 0 \quad (55)$$

So finally J10 is given by

$$\left( \frac{6}{\beta^2} D_x^{10} + \frac{18}{\beta} D_x^5 D_t + D_t^2 + c_1 D_x^3 (D_\tau + D_x^7) + c_2 D_x^3 (D_{\tau'} + D_x^2 D_t) \right) f \cdot f = 0 \quad (56)$$

$$D_x (D_\tau + D_x^7) f \cdot f = 0 \quad (57)$$

$$D_x (D_{\tau'} + D_x^2 D_t) f \cdot f = 0 \quad (58)$$

Summarizing all the above, such equations can be considered as the  $l$ -linear dominant part of order  $l + m - 1$  parametrized by bilinear relations.

$$D_x^l(D_\tau + D_x^m)(f \cdot f) + \dots = \text{the equation}, \quad (59)$$

$$D_x(D_\tau + D_x^m)(f \cdot f) = 0 \quad (60)$$

We will denote it as  $(D^l, D^m)$ . The following table represents  $l, m$  bilinear pairs of lowest orders:

| $m, l$ | 3      | 5    | 7     | 9 | 11 |
|--------|--------|------|-------|---|----|
| 3      | Lax5   | Lax7 | -     | - | -  |
| 5      | SK7/H8 | H10  | -     | - | -  |
| 7      | J10    | -    | new13 | - | -  |
| 9      | -      | -    | -     | - | -  |
| 11     | -      | -    | -     | - | -  |

## 5. Higher order equations

In this section we represent some results on higher order generalizations of J8, J10 equations. Firstly, we formulate the principle.

**Property. 1.** For  $D^n$  Hirota operator fuchs array is symmetric over negative indices

**Property. 2.** If equations  $E_1, E_2, \dots$  are in a KP-type hierarchy with  $\deg(E_1) < \deg(E_2) < \dots$ , then

$$s_1 \subset s_2 \subset \dots, \quad (61)$$

where  $s_i$  is a triangular sub-array of negative fuchs indices of  $E_i$

Using these properties, we can build higher order analogues of J10 equation. The equations should be organized in the way (in a manner similar to the KP, BKP hierarchies [6]):

$$\begin{aligned} D^{10} + (D^3, D^7) &= J10, \\ D^{12} + (D^5, D^7) + (D^3, J10) &= J12, \\ D^{14} + (D^7, D^7) + (D^5, J10) + (D^3, J12) &= J14, \\ &\dots \end{aligned}$$

Namely, consider degree-homogeneous linear combination (see 59)

$$D^{12} + k_1(D^5, D^7) + k_2(D^3, J10) \quad (62)$$

We can obtain the coefficients  $k_i$  resolving all fuchs indices one by one, so that

$$F = \begin{pmatrix} -1.0 & 1.0 & 2.0 & 3.0 & 4.0 & \dots \\ -2.0 & -1.0 & 1.0 & 2.0 & 3.0 & \\ -3.0 & -2.0 & -1.0 & 1.0 & 4.0 & \dots \\ -13.0 & -11.0 & -7.0 & -1.0 & 0.36985419 & \dots \\ -13.76327806 & -11.56856256 & -7.247114834 & -0.3858031569 & -1.0 & \dots \end{pmatrix} \quad (63)$$

The corresponding values are

$$k_1 = 27, k_2 = -28. \quad (64)$$

However, due to the absence of irrational indices, this equation might not be integrable. But it is closely related to the determinat-like 12th order extension of J10 (65), which is already integrable!

$$\begin{aligned} J12 = & -18 P_{[1,0,5,6]} + 4 P_{[1,5,0,6]} + 5 P_{[1,10,0,1]} - 80 P_{[3,3,3,3]} - 35 P_{[3,4,1,4]} + \\ & + 10 P_{[4,0,1,7]} + 60 P_{[4,2,2,4]} + 30 P_{[4,3,0,5]} + 24 P_{[5,1,1,5]} \end{aligned} \quad (65)$$

Expanding all pentilinear operators in 65, we get:

$$\begin{aligned} J12 = & \left( 2 f_{8x} f_{4x} - 2 f_{7x} f_{5x} - 2 f_{9x} f_{3x} + f_{6x}^2 + f_{10x} f_{2x} \right) f^3 + \left( 28 f_{6x} f_{3x}^2 - f_{10x} f_x^2 + 9 f_{8x} f_{2x}^2 + \right. \\ & + 2 f_{5x} f_{6x} f_x + 6 f_{5x}^2 f_{2x} + 10 f_{8x} f_x f_{3x} - 4 f_{9x} f_x f_{2x} - 30 f_{5x} f_{3x} f_{4x} - 6 f_{7x} f_{4x} f_x + \\ & \left. + 10 f_{4x}^3 + 4 f_{6x} f_{2x} f_{4x} - 28 f_{7x} f_{2x} f_{3x} \right) f^2 + \dots \end{aligned}$$

## 6. Summary

As we have shown, multilinear Hirota equations arise from the KP-type bilinear hierarchies, i.e. they can be expanded as  $n$ -tuples of bilinear equations, each of them is contained in a hierarchy (we recall here that such hierarchies are intimately linked with the affine Kac-Moody algebras). This fact follows from the Lie-algebraic structure of these equations. Thus, gauge-invariant multilinear operators of higher order can be useful for describing such higher reductions. However, when the reduction is done, total dimension is not greater than  $(2+1)$ . It is also an interesting fact that J10 integrable equation consists of 3 bilinear equations, one of which is not integrable in the “default” time-variables. But its  $(2+1)$  extension is truly integrable (can be obtained by adding BKP even terms to J10 and assuming a solution in form of diagonal Schur function). We also find J12 analogue as the determinant-like equation (it is non-evolutionary and dispersionless). And it remains unclear if there are connections with hyperbolic Kac-Moody algebras. The reasonable approach is as follows. One should extend the space of symmetric functions (using some natural deformations) and define an operator similar to Hirota derivative, which eigenvectors are connected with  $U_q(\mathfrak{sl}_2(\mathbb{C}))$ . This is of further development.

## References

- [1] C. Athorne Algebraic invariants and generalized Hirota derivatives, Physics Letters A, vol. 256 n. 1 p. 20, 1999
- [2] M. England, C. Athorne Building Abelian Functions with Generalised BakerHirota Operators, SIGMA 8, 2012, 037
- [3] A. Goriely Integrability and Nonintegrability of Dynamical Systems, Advanced Series in Nonlinear Dynamics vol. 19, WSP, 2001

- [4] J. Hietarinta, B. Grammaticos Integrable trilinear PDE's, arXiv preprint solv-int/9411003, 1994
- [5] J. Hietarinta Hirota's bilinear method and soliton solutions, Physics AUC, vol. 15 (part I), 31-37, 2005
- [6] M. Jimbo, T. Miwa Solitons and Infinite Dimensional Lie Algebras, Publ RIMS, Kyoto Univ. 19, 1983, 943-1001
- [7] S. Lu On Soliton Equations of Exceptional Type, Journal of Algebra 166, 611-629, 1994